

## An Updated View of Breast Reconstruction

Yuri Jiménez Caprielova<sup>1</sup>, Kenia Selene Avilés Palomares<sup>2</sup>, Dr. Juan Carlos Hernández Torón<sup>3</sup>

<sup>1,2</sup>Residente de Cirugía Plástica y Reconstructiva, IMSS Hospital de Traumatología y Ortopedia "Dr. Victorio de la Fuente Narváez"

<sup>3</sup>Titular y Adscrito de cirugía plástica y reconstructiva, IMSS Hospital de Traumatología y Ortopedia "Dr. Victorio de la Fuente Narváez"

### ABSTRACT

Breast reconstruction is a cornerstone of post-mastectomy care, aiming to restore the breast's form and appearance while enhancing patients' psychosocial well-being. The procedure can be performed using implant-based or autologous tissue techniques, each tailored to the patient's medical profile, treatment plan, and preferences. Implant-based reconstruction, involving saline or silicone implants, offers shorter recovery times and is often preferred in patients without planned postmastectomy radiation therapy. Autologous tissue reconstruction, including techniques like the DIEP and latissimus dorsi flaps, provides natural contours and greater durability, particularly beneficial for patients undergoing radiation therapy. Advancements such as nipple-sparing mastectomy, acellular dermal matrices, and minimally invasive approaches have improved surgical outcomes and patient satisfaction. However, complications—ranging from implant-related issues like capsular contracture to donor-site morbidity—highlight the need for comprehensive preoperative planning and a multidisciplinary approach. Breast reconstruction remains a highly individualized process, emphasizing patient-centered care to optimize oncological safety, aesthetic outcomes, and quality of life.

**KEYWORDS:** Breast reconstruction, mastectomy, implant-based reconstruction, autologous tissue reconstruction, DIEP flap, latissimus dorsi flap, postmastectomy radiation therapy, nipple-sparing mastectomy, patient-centered care, reconstructive surgery.

### ARTICLE DETAILS

**Published On:**  
**19 December 2024**

**Available on:**  
<https://ijmscr.org/>

### INTRODUCTION

Breast reconstruction is a surgical procedure aimed at restoring the shape and appearance of a breast following mastectomy or breast-conserving surgery. It is an integral part of breast cancer treatment for many patients, offering both aesthetic and psychosocial benefits.<sup>[1-3]</sup> The reconstruction can be performed immediately during the mastectomy or in a delayed fashion, depending on the patient's medical condition, treatment plan, and personal preferences.<sup>[1][4]</sup>

There are two primary types of breast reconstruction: implant-based and autologous tissue reconstruction. Implant-based reconstruction involves the use of saline or silicone implants, which can be placed immediately or after a tissue expander has been used to prepare the site.<sup>[1][5]</sup> Autologous tissue reconstruction uses the patient's own tissue, often from the abdomen, back, or buttocks, to create a new breast mound. Techniques such as the transverse rectus abdominis myocutaneous (TRAM) flap, latissimus dorsi flap, and deep

inferior epigastric perforator (DIEP) flap are commonly used.<sup>[1-2][5]</sup>

Advancements in surgical techniques, such as nipple-sparing mastectomy and the use of acellular dermal matrices, have improved the aesthetic outcomes of breast reconstruction.<sup>[3]</sup> The choice of reconstruction method depends on various factors, including the patient's body habitus, smoking history, comorbidities, and whether postmastectomy radiation therapy is planned, as radiation can affect the outcomes of reconstruction.<sup>[1][4]</sup>

Overall, breast reconstruction is a personalized decision that should be made in consultation with a multidisciplinary team, taking into account the patient's medical needs and personal preferences.<sup>[1-2][4]</sup>

#### Indications

Breast reconstruction surgery following mastectomy or breast-conserving surgery is primarily indicated to restore breast symmetry and improve the patient's quality of life by

## An Updated View of Breast Reconstruction

addressing cosmetic, body image, and psychosocial concerns. The choice between implant-based and autologous tissue reconstruction depends on several factors, including the patient's medical condition, treatment plan, and personal preferences.

Implant-based reconstruction is often chosen for its shorter operative time and quicker recovery, making it suitable for patients who prefer a less invasive procedure. It involves either a single-stage reconstruction with a permanent implant or a two-stage process with tissue expansion followed by implant placement. This method is generally preferred when post-mastectomy radiation therapy (PMRT) is not planned, as radiation can increase the risk of complications such as capsular contracture.<sup>[6-7]</sup>

Autologous tissue reconstruction, which uses the patient's own tissue, is often considered when PMRT is indicated, as it tends to provide more stable long-term results despite the complexity of the procedure. Common donor sites include the abdomen, back, and buttocks, with techniques such as the TRAM flap or latissimus dorsi flap being utilized. This method is advantageous in patients with sufficient donor tissue and those who may have contraindications to implants, such as a history of smoking or diabetes, which can increase complication rates.<sup>[7-8]</sup>

Patient preference plays a crucial role in the decision-making process, as some may prioritize a more natural feel and appearance, which autologous reconstruction can offer, while others may opt for the simplicity and shorter recovery associated with implants. Additionally, the timing of reconstruction (immediate vs. delayed) is influenced by the overall treatment plan, including the need for adjuvant therapies like radiation.<sup>[9-11]</sup>

Ultimately, a multidisciplinary approach involving oncologists, plastic surgeons, and the patient is essential to tailor the reconstructive strategy to the individual's needs and circumstances, ensuring both oncological safety and aesthetic satisfaction.<sup>[7][9]</sup>

### Complications

Breast reconstruction surgery, while beneficial, is associated with several potential complications, which can vary based on factors such as the type of reconstruction, patient medical history, and the use of postmastectomy radiation therapy (PMRT).

#### 1. Implant-Based Reconstruction Complications:

Common complications include infection, capsular contracture, and implant loss. The use of PMRT significantly increases the risk of complications such as infection and implant removal. For instance, radiation therapy has been associated with higher odds of implant removal in patients with implant-based reconstruction. Additionally, larger implant sizes and factors like low plasma albumin and elevated blood glucose levels can increase the risk of complications.<sup>[12-14]</sup>

**2. Autologous Tissue Reconstruction Complications:** This method can lead to complications such as fat necrosis, flap loss, and donor site morbidity. PMRT can exacerbate these issues, leading to parenchymal complications like fat necrosis and fibrosis, as well as skin complications such as tissue retraction and hypertrophic scarring. Smoking and diabetes are significant risk factors for these complications.<sup>[15]</sup>

**3. General Complications Across Techniques:** Both reconstruction types can experience complications such as seroma, hematoma, and wound dehiscence. Factors like higher body mass index (BMI), smoking, and diabetes increase the risk of these complications. Nipple-sparing mastectomy, while generally safe, can have specific complications like nipple necrosis, especially in patients with preoperative irradiation.<sup>[16-17]</sup>

**4. Impact of Patient Factors:** Patient-specific factors such as age, BMI, and comorbidities like diabetes and smoking status significantly influence complication rates. For example, higher BMI and smoking are associated with increased risks of skin necrosis and other complications.<sup>[16-17]</sup>



Figure 1. Reconstruction with tissue expanders



Figure 2. Immediate PO period with tissue expanders

#### Latissimus Dorsi Flap reconstruction

The latissimus dorsi flap (LDF) technique is a well-established method for breast reconstruction, offering several considerations and outcomes that are important for clinical decision-making.

The LDF is versatile and can be used for both immediate and delayed breast reconstruction, often in combination with implants to achieve the desired breast volume, especially in patients with larger breasts.<sup>[18-19]</sup> It is particularly valuable in

## An Updated View of Breast Reconstruction

cases where other reconstruction options, such as abdominal flaps, are not feasible or when patients have undergone radiation therapy, as the autologous tissue can help mitigate some of the adverse effects of radiation.<sup>[18][20]</sup>

Outcomes associated with the LDF technique are generally favorable. Studies have shown high patient satisfaction rates, with many patients reporting good aesthetic outcomes and psychosocial well-being.<sup>[21-22]</sup> The technique is associated with a relatively low rate of serious complications. Common complications include seroma formation, which is the most frequently reported issue, occurring in up to 26% of cases.<sup>[18-19]</sup> Other complications such as wound dehiscence, infection, and flap necrosis are less common.<sup>[18-19][22]</sup>

Donor-site morbidity, particularly related to shoulder function, is a consideration with the LDF technique. Some studies have reported a decrease in shoulder strength postoperatively, although this does not typically impair activities of daily living.<sup>[22]</sup> The use of immediate fat transfer with the LDF, known as the LIFT technique, has been explored to enhance volume and aesthetic outcomes, although it often requires additional fat grafting procedures.<sup>[23]</sup>

Overall, the LDF remains a reliable and safe option for breast reconstruction, particularly in specific clinical scenarios where other methods may not be suitable. The choice of technique should be individualized based on patient characteristics, preferences, and clinical circumstances.<sup>[18][24]</sup>



Figure 3. Donor site



Figure 4. Postoperative image

### MASTOPEXY

Mastopexy, in the context of breast reconstruction, is a surgical procedure aimed at lifting and reshaping ptotic (sagging) breasts, often performed in conjunction with other

reconstructive techniques to optimize aesthetic outcomes. This procedure can be particularly beneficial for patients with large or ptotic breasts undergoing nipple-sparing mastectomy (NSM) and reconstruction, as it helps manage the redundant skin envelope and reduces the risk of ischemic complications.<sup>[25-26]</sup>

In comparison to the latissimus dorsi flap technique, mastopexy serves a different purpose. The latissimus dorsi flap is an autologous tissue reconstruction method that provides additional tissue coverage, which can be advantageous for patients who have undergone radiation therapy, as it may improve the vascularity and healing of the reconstructed breast. This technique is often chosen for its reliability and the ability to provide a natural contour, especially in patients with a history of radiation, where tissue quality is compromised.

For patients with larger breasts or those who have undergone radiation therapy, a staged approach involving mastopexy before NSM and reconstruction can be advantageous. This approach has been shown to decrease the risk of complications and improve clinical outcomes by preparing the breast for subsequent reconstructive procedures, such as implant-based reconstruction or the use of free abdominal flaps.<sup>[25-26]</sup> In contrast, the latissimus dorsi flap may be more suitable for patients requiring additional tissue coverage and those with compromised skin quality due to radiation.

Ultimately, the choice between mastopexy and techniques like the latissimus dorsi flap depends on individual patient factors, including breast size, degree of ptosis, history of radiation therapy, and personal preferences. Each technique has its own set of indications and potential complications, and the decision should be tailored to achieve the best possible aesthetic and functional outcomes for the patient.

### DIEP

The deep inferior epigastric perforator (DIEP) flap technique is a widely used method in breast reconstruction that involves the transfer of skin and fat from the lower abdomen to the chest to reconstruct a breast following mastectomy. This technique is muscle-sparing, as it utilizes the perforating branches of the deep inferior epigastric artery (DIEA) without sacrificing the rectus abdominis muscle, thereby reducing donor site morbidity compared to traditional methods like the transverse rectus abdominis myocutaneous (TRAM) flap.<sup>[27]</sup>

The DIEP flap is advantageous because it preserves abdominal muscle function, which decreases the risk of complications such as abdominal wall weakness or hernia. Preoperative imaging, such as computed tomography angiography, is often employed to map the vascular anatomy and identify the most suitable perforators, which can vary significantly among patients.<sup>[11]</sup> This imaging helps optimize the surgical plan, potentially reducing operative time and improving outcomes.

Recent advancements in the DIEP flap technique include minimally invasive approaches, such as laparoscopic and robotic-assisted dissections, which aim to further minimize

## An Updated View of Breast Reconstruction

donor site trauma and improve recovery times. These approaches have shown promise in reducing the length of fascial incisions and associated morbidity.<sup>[28-30]</sup> Additionally, techniques like the short-fasciotomy method have been developed to limit the extent of fascial dissection, thereby preserving more of the abdominal wall structure and function.<sup>[31]</sup>

Overall, the DIEP flap is a robust option for autologous breast reconstruction, offering aesthetic and functional benefits while minimizing donor site complications. The choice of technique and approach can be tailored to the patient's specific anatomy and clinical needs, guided by preoperative planning and imaging.<sup>[28-31]</sup>



**Figure 5. DIEP reconstruction**

### TRAM

The transverse rectus abdominis myocutaneous (TRAM) flap is a well-established technique in breast reconstruction that utilizes tissue from the lower abdomen to create a new breast mound. This method can be performed as either a pedicled or free flap. The pedicled TRAM flap is based on the superior epigastric vessels and involves tunneling the flap under the skin to the chest, while the free TRAM flap involves complete detachment and microvascular anastomosis to the chest vessels, typically the thoracodorsal or internal mammary vessels.<sup>[32-33]</sup>

In comparison to the latissimus dorsi flap and the deep inferior epigastric perforator (DIEP) flap, the TRAM flap has distinct indications and patient profiles. The latissimus dorsi flap, which uses muscle and skin from the back, is often chosen when additional tissue coverage is needed, particularly in patients with compromised skin quality due to radiation therapy. It is less commonly used as a standalone option for breast volume but can be combined with implants.<sup>[34]</sup>

The DIEP flap, on the other hand, is a muscle-sparing technique that uses only skin and fat from the abdomen, preserving the rectus abdominis muscle. This approach reduces donor site morbidity, such as abdominal wall weakness and hernia, which are more common with the TRAM flap due to muscle sacrifice.<sup>[33-34]</sup> The DIEP flap is particularly suitable for patients who prioritize minimizing

abdominal complications and for those with adequate perforator vessels, as it requires meticulous surgical technique and preoperative planning.<sup>[35-36]</sup>

The TRAM flap remains a viable option, especially in settings where microsurgical expertise for DIEP flaps is not available or in patients who may not be ideal candidates for the DIEP flap due to vascular anatomy or other factors. It is often used in patients without significant risk factors for flap loss, such as smoking or obesity, although modifications like the free TRAM can be employed in higher-risk patients to improve outcomes.<sup>[36-37]</sup>



**Figure 6. Preoperative image**



**Figure 7. Postoperative image for TRAM flap reconstruction**

### CONCLUSION

Breast reconstruction is a vital component of post-mastectomy care, offering numerous aesthetic and psychosocial benefits for patients recovering from breast cancer treatment. The choice of reconstruction technique—whether implant-based or autologous tissue—requires a highly individualized approach, considering factors such as the patient's medical condition, treatment plan, and personal preferences. Advanced techniques like the DIEP flap, latissimus dorsi flap, and nipple-sparing mastectomy continue to enhance outcomes, improving both functionality and appearance.

While complications, including capsular contracture, flap loss, and donor-site morbidity, can occur, meticulous surgical planning, preoperative imaging, and multidisciplinary

## An Updated View of Breast Reconstruction

collaboration help minimize risks and optimize results. Each reconstructive option, from implant-based methods to autologous tissue techniques, has unique advantages and potential drawbacks, making shared decision-making crucial in tailoring care to the individual patient.

Ultimately, breast reconstruction represents not just a surgical procedure but a pathway to restoring confidence and quality of life, underscoring the importance of personalized, patient-centered care in achieving the best outcomes.

### REFERENCES

- I. Invasive Breast Cancer. Carlson RW, Allred DC, Anderson BO, et al. *Journal of the National Comprehensive Cancer Network: JNCCN*. 2011;9(2):136-222. doi:10.6004/jnccn.2011.0016. *Leading Journal*
- II. Update on Breast Reconstruction Techniques and Indications. Petit JY, Rietjens M, Lohsiriwat V, et al. *World Journal of Surgery*. 2012;36(7):1486-97. doi:10.1007/s00268-012-1486-3.
- III. Breast Reconstruction. Sbitany H. *The Surgical Clinics of North America*. 2018;98(4):845-857. doi:10.1016/j.suc.2018.03.011.
- IV. Breast Reconstruction After Mastectomy. Tondu T, Tjalma WAA, Thiessen FEF. *European Journal of Obstetrics, Gynecology, and Reproductive Biology*. 2018;230:228-232. doi:10.1016/j.ejogrb.2018.04.016.
- V. ACR Appropriateness Criteria® Imaging After Mastectomy and Breast Reconstruction. Heller SL, Lourenco AP, Niell BL, et al. *Journal of the American College of Radiology : JACR*. 2020;17(11S):S403-S414. doi:10.1016/j.jacr.2020.09.009.
- VI. Breast Reconstruction With Prosthetic Implants. Mesbahi AN, McCarthy CM, Disa JJ. *Cancer Journal (Sudbury, Mass.)*. 2008 Jul-Aug;14(4):230-5. doi:10.1097/PPO.0b013e31817fb7c3.
- VII. Breast Reconstruction Following Cancer Treatment. Gerber B, Marx M, Untch M, Faridi A. *Deutsches Arzteblatt International*. 2015;112(35-36):593-600. doi:10.3238/arztebl.2015.0593.
- VIII. Invasive Breast Cancer. Carlson RW, Allred DC, Anderson BO, et al. *Journal of the National Comprehensive Cancer Network : JNCCN*. 2011;9(2):136-222. doi:10.6004/jnccn.2011.0016.
- IX. Breast Reconstruction Options Following Mastectomy. Djohan R, Gage E, Bernard S. *Cleveland Clinic Journal of Medicine*. 2008;75 Suppl 1:S17-23. doi:10.3949/ccjm.75.suppl\_1.s17.
- X. Update on Breast Reconstruction Techniques and Indications. Petit JY, Rietjens M, Lohsiriwat V, et al. *World Journal of Surgery*. 2012;36(7):1486-97. doi:10.1007/s00268-012-1486-3.
- XI. Current Trends in Postmastectomy Breast Reconstruction. Panchal H, Matros E. *Plastic and Reconstructive Surgery*. 2017;140(5S Advances in Breast Reconstruction):7S-13S. doi:10.1097/PRS.0000000000003941.
- XII. Complications After Mastectomy and Immediate Breast Reconstruction for Breast Cancer: A Claims-Based Analysis. Jagsi R, Jiang J, Momoh AO, et al. *Annals of Surgery*. 2016;263(2):219-27. doi:10.1097/SLA.0000000000001177.
- XIII. Impact of Radiotherapy on Complications and Patient-Reported Outcomes After Breast Reconstruction. Jagsi R, Momoh AO, Qi J, et al. *Journal of the National Cancer Institute*. 2018;110(2). doi:10.1093/jnci/djx148.
- XIV. Single-Institution Algorithm for Prevention and Management of Complications in Direct-to-Implant Breast Reconstruction. Yin Z, Wang H, Liu Y, et al. *Plastic and Reconstructive Surgery*. 2022;150:48S-60S. doi:10.1097/PRS.0000000000009490.
- XV. Irradiated Autologous Breast Reconstructions: Effects of Patient Factors and Treatment Variables. Albino FP, Koltz PF, Ling MN, Langstein HN. *Plastic and Reconstructive Surgery*. 2010;126(1):12-16. doi:10.1097/PRS.0b013e3181da878f.
- XVI. Breast Reconstruction Following Nipple-Sparing Mastectomy: Predictors of Complications, Reconstruction Outcomes, and 5-Year Trends. Colwell AS, Tessler O, Lin AM, et al. *Plastic and Reconstructive Surgery*. 2014;133(3):496-506. doi:10.1097/01.prs.0000438056.67375.75.
- XVII. Complications in Postmastectomy Breast Reconstruction: One-Year Outcomes of the Mastectomy Reconstruction Outcomes Consortium (MROC) Study. Wilkins EG, Hamill JB, Kim HM, et al. *Annals of Surgery*. 2018;267(1):164-170. doi:10.1097/SLA.0000000000002033.
- XVIII. Latissimus Dorsi Flap for Breast Reconstruction: A Large Single-Institution Evaluation of Surgical Outcome and Complications. Banyas-Paluchowski M, Brus L, Krawczyk N, et al. *Archives of Gynecology and Obstetrics*. 2024;309(1):269-280. doi:10.1007/s00404-023-07186-3.
- XIX. Comparison of Different Techniques in Latissimus Dorsi Breast Reconstruction: Plain, Immediately Lipofilled, and Implant Enhanced. Palve J, Luukkaala T, Kääriäinen M. *Annals of Plastic Surgery*. 2022;88(1):20-24. doi:10.1097/SAP.0000000000002798.
- XX. Immediate Latissimus Dorsi and Prosthetic Reconstruction in the Setting of Postmastectomy Radiation: An Analysis of 376 Breast Reconstructions. Chiasson KF, Kumbala PA, Restrepo RD, Soto E, Cohn AB. *Annals of Plastic*

## An Updated View of Breast Reconstruction

- Surgery. 2020;84(6S Suppl 5):S364-S368. doi:10.1097/SAP.0000000000002279.
- XXI. Long-Term Outcomes of Latissimus Dorsi Flap Breast Reconstructions: A Single-Centre Observational Cohort Study With Up to 12 Years of Follow Up. Wattoo G, Nayak S, Khan S, et al. *Journal of Plastic, Reconstructive & Aesthetic Surgery : JPRAS.* 2021;74(9):2202-2209. doi:10.1016/j.bjps.2020.12.058.
- XXII. A Retrospective Study of Latissimus Dorsi Flap in Immediate Breast Reconstruction. Zheng H, Zhu G, Guan Q, et al. *Frontiers in Oncology.* 2021;11:598604. doi:10.3389/fonc.2021.598604.
- XXIII. Donor-Site Morbidity Following Breast Reconstruction With a Latissimus Dorsi Flap - A Prospective Study. Højvig JH, Henriksen M, Bartholdy CR, Bonde CT. *Journal of Plastic, Reconstructive & Aesthetic Surgery : JPRAS.* 2022;75(7):2205-2210. doi:10.1016/j.bjps.2022.01.036.
- XXIV. Breast Reconstruction Using the Latissimus Dorsi Flap and Immediate Fat Transfer (LIFT): A Systematic Review and Meta-Analysis. Escandón JM, Escandón L, Ahmed A, et al. *Journal of Plastic, Reconstructive & Aesthetic Surgery : JPRAS.* 2022;75(11):4106-4116. doi:10.1016/j.bjps.2022.08.025.
- XXV. Staged Mastopexy Before Nipple-Sparing Mastectomy: Improving Safety and Appearance in Breast Reconstruction. Awaida CJ, Bernier C, Bou-Merhi JS, et al. *Plastic and Reconstructive Surgery.* 2024;153(5):864e-872e. doi:10.1097/PRS.0000000000010823.
- XXVI. The Role of Premastectomy Mastopexy and Breast Reduction in the Reconstruction of the Enlarged or Ptotic Breast. Hammond DC, Little AK. *Plastic and Reconstructive Surgery.* 2022;150(2):270-280. doi:10.1097/PRS.0000000000009340.
- XXVII. ACR Appropriateness Criteria® Imaging of Deep Inferior Epigastric Arteries for Surgical Planning (Breast Reconstruction Surgery): 2022 Update. Singh N, Aghayev A, Ahmad S, et al. *Journal of the American College of Radiology : JACR.* 2022;19(11S):S357-S363. doi:10.1016/j.jacr.2022.09.004.
- XXVIII. Minimally Invasive Laparoscopically Dissected Deep Inferior Epigastric Artery Perforator Flap: An Anatomical Feasibility Study and a First Clinical Case. Hivelin M, Soprani A, Schaffer N, Hans S, Lantieri L. *Plastic and Reconstructive Surgery.* 2018;141(1):33-39. doi:10.1097/PRS.0000000000003989.
- XXIX. Robotic Harvest of the Deep Inferior Epigastric Perforator Flap for Breast Reconstruction: A Case Series. Bishop SN, Asaad M, Liu J, et al. *Plastic and Reconstructive Surgery.* 2022;149(5):1073-1077. doi:10.1097/PRS.0000000000008988.
- XXX. Robotic DIEP Flap Harvest Through a Totally Extraperitoneal Approach Using a Single-Port Surgical Robotic System. Choi JH, Song SY, Park HS, et al. *Plastic and Reconstructive Surgery.* 2021;148(2):304-307. doi:10.1097/PRS.0000000000008181.
- XXXI. Short Fasciotomy-Deep Inferior Epigastric Perforator Flap Harvest for Breast Reconstruction. Kim J, Lee KT, Mun GH. *Plastic and Reconstructive Surgery.* 2023;152(6):972e-984e. doi:10.1097/PRS.0000000000010382.
- XXXII. Breast Reconstruction With the TRAM Flap: Pedicled and Free. Serletti JM. *Journal of Surgical Oncology.* 2006;94(6):532-7. doi:10.1002/jso.20492.
- XXXIII. Optimizing Pedicled Transverse Rectus Abdominis Muscle Flap Breast Reconstruction. Kanchwala SK, Bucky LP. *Cancer Journal (Sudbury, Mass.).* 2008 Jul-Aug;14(4):236-40. doi:10.1097/PPO.0b013e318180bce5.
- XXXIV. Choice of Flaps for Breast Reconstruction. Tachi M, Yamada A. *International Journal of Clinical Oncology.* 2005;10(5):289-97. doi:10.1007/s10147-005-0527-4.
- XXXV. A Comparison Study of Deep Muscle Sparing Transverse Rectus Abdominis Musculocutaneous Flap for Breast Reconstruction. Tokumoto H, Akita S, Arai M, et al. *Microsurgery.* 2019;39(7):583-589. doi:10.1002/micr.30443.
- XXXVI. Muscle Sparing-2 Transverse Rectus Abdominis Musculocutaneous Flap for Breast Reconstruction: A Comparison With Deep Inferior Epigastric Perforator Flap. Takeishi M, Fujimoto M, Ishida K, Makino Y. *Microsurgery.* 2008;28(8):650-5. doi:10.1002/micr.20563.
- XXXVII. A Safe and Efficient Technique for Pedicled TRAM Flap Breast Reconstruction. Semple JL, Viesel-Mathieu A, Alshaqsi S, Armstrong K. *Plastic and Reconstructive Surgery.* 2023;151(6):1022e-1034e. doi:10.1097/PRS.0000000000010174.