

Liposuction and Body Contouring Surgery: A Comprehensive Review of Techniques, Safety, And Evolving Evidence

Ivan Yahir Paternina González^{*1}, Abraham Saynes Jarquin², Cambises Juda Bernal Fausto³, Irvin Hernández Sánchez⁴, Gilberto Luis Juan Baltazar⁵, Marco Antonio Hernández Hernández¹, César Iván García Calderón⁶, Diego Gabriel Espinosa Huerta³, Ethel Cassandra Ramírez Ortiz⁷, Clyvia Coral Medina Lavenant⁸, Alejandro Ramirez Rodeles⁹, Nicole Rentería Vázquez¹⁰, Cristobal Montes De Oca Fabila¹¹.

¹Hospital Español de México.

²HGZ 48, IMSS.

³UMAE Hospital de Especialidades Centro Médico Nacional Siglo XXI “Dr Bernardo Sepúlveda Gutiérrez”.

⁴Hospital General de México “Dr. Eduardo Liceaga”.

⁵Centro universitario del sur Universidad de Guadalajara.

⁶IMSS Hospital General Regional #1.

⁷Hospital Ángeles, CDMX.

⁸Hospital general de zona 16 IMSS.

⁹Hospital General Regional Numero 1 IMSS. Ciudad Obregon

¹⁰Universidad de Guadalajara (Cusur).

¹¹Hospital General del Estado de Sonora Dr. Ernesto Ramos Bours.

ABSTRACT

Liposuction remains among the most frequently performed cosmetic surgical procedures worldwide and continues to evolve through technological refinement and accumulating clinical evidence. This review synthesizes contemporary literature on the history, anatomical basis, techniques, anesthetic principles, complication profile, and emerging adjuncts of suction-assisted body contouring. Traditional suction-assisted lipectomy (SAL) remains the most widely used technique, but energy-based modalities—ultrasound-assisted liposuction (UAL), power-assisted lipectomy (PAL), laser-assisted lipectomy (LAL), radiofrequency-assisted lipectomy (RFAL), and water-assisted lipectomy (WAL)—have been developed to reduce operator fatigue, decrease blood loss, and, in some cases, promote skin retraction. Tumescence anesthesia, introduced by Klein, transformed liposuction safety by enabling large-volume fat removal under local anesthesia with minimal blood loss. Reported complication rates remain low overall, with contour deformity the most common adverse outcome and gluteal fat transfer carrying the highest procedure-specific mortality risk among aesthetic interventions, primarily related to fat embolism. Comparative data on ultrasound-assisted liposuction suggest a consistent reduction in intraoperative blood loss relative to traditional suction-assisted techniques, although evidence for superior long-term skin tightening remains inconclusive. Pharmacologic adjuncts such as tranexamic acid have shown mixed results across studies, with effectiveness appearing dependent on dose and route of administration. Noncosmetic indications, including lipedema, lipodystrophy, and gynecomastia, continue to expand the clinical utility of the procedure. Overall, liposuction maintains a favorable safety profile when performed by trained surgeons in appropriately selected patients, though continued investigation is warranted to clarify the comparative advantages of newer energy-based technologies.

KEYWORDS: Ultrasound-assisted liposuction, laser-assisted lipectomy, power-assisted lipectomy, traditional suction-assisted lipectomy.

ARTICLE DETAILS

Published On:
15 July 2026

Available on:
<https://ijmscrs.com>

I. INTRODUCTION

Suction-assisted lipectomy, more commonly known as liposuction, is an outpatient procedure designed to remove adipose tissue from the subcutaneous space in order to achieve a more desirable body contour (1). It is the second most performed cosmetic surgery in the United States and the most common surgical procedure among patients between the ages of 35 and 64 (1). In 2018, surgeons performed 258,558 liposuction procedures, representing a 5 percent increase from the prior year, and the overall number of liposuction procedures increased 124 percent between 1997 and 2015 (1). More recent global estimates place the number of liposuction procedures performed worldwide at approximately 300,000 annually, with the American Society of Plastic Surgeons reporting 2.3 million total plastic surgery procedures performed globally in 2020 (2).

Beyond its purely aesthetic applications, liposuction serves as an important adjunct in reconstructive surgery, particularly when harvested fat is autologously reinjected into tissues of the breast or face (1). One procedure that has risen sharply in popularity and drawn considerable media attention is gluteal fat transfer, commonly known as the “Brazilian butt lift” (1). At the same time, the rapid proliferation of assistive technologies—ultrasonic, powered, laser-based, and radiofrequency devices—has generated debate over whether such innovations meaningfully improve outcomes or primarily serve as marketing differentiators with limited proven clinical benefit (3). This review integrates findings from historical accounts, original clinical investigations, and systematic literature analyses to provide an updated, evidence-based overview of liposuction: its history, anatomical principles, indications, techniques, anesthetic considerations, complications, and the comparative evidence surrounding blood loss and adjunctive pharmacologic strategies such as tranexamic acid.

II. HISTORICAL DEVELOPMENT

The history of liposuction is marked by a slow and often hazardous evolution toward the safer, more controlled techniques used today. The first documented attempt at surgical fat removal was performed by Charles Dujarrier in 1921, who used a uterine curette to operate on the knees and calves of a dancer; the procedure resulted in injury to the femoral artery and ultimately required amputation of the limb (1). This tragic outcome led to the near-abandonment of fat-removal surgery for several decades. In 1964, Josef Schrudde attempted a similar curettage technique on a patient's leg but observed skin necrosis in four of fifteen

treated patients, along with hematoma and seroma formation (1).

The modern era of liposuction began in the mid-1970s. In 1974, the father-son team of Giorgio and Arpad Fischer attempted fat removal using a hollow cannula (1), and by 1975 they had pioneered the use of blunt hollow cannulas combined with suction curettage for liposuction of the outer thighs, although patients in this series experienced deforming lymphorrhea (1). It was French dermatologist Yves-Gerard Illouz, however, who is widely credited with popularizing modern liposuction in the late 1970s through the combination of a hollow, blunt-tipped cannula with suction, thereby eliminating the dangers associated with sharp curettes (1). In 1977, Illouz also developed the “wet technique,” in which hypotonic saline solution and hyaluronidase were injected into adipose tissue prior to liposuction to reduce hemorrhagic risk—a form of hydrodissection that preserved neurovascular bundles and facilitated aspiration (1).

Subsequent refinements followed in rapid succession. In 1983, Pierre Fournier introduced the use of syringes rather than mechanical suction to allow for finer control of negative pressure, and also contributed the “criss-cross” technique of cannula movement (1,5,6). Luiz Toledo expanded the use of syringes to multiple gauges and sizes for fat aspiration in 1988 (1). The single most transformative advance, however, came in 1987, when American dermatologist Jeffrey Klein introduced the tumescent technique—a method of local anesthesia infiltration using dilute lidocaine and epinephrine that permitted removal of larger fat volumes while substantially reducing bleeding (1,5,6). Because 30 to 45 percent of lipoaspirate volume consists of blood when liposuction is performed without prior fluid instillation, the dry technique used in early liposuction has been almost entirely abandoned in favor of tumescent or superwet approaches (1).

Technological refinement continued into the 1990s. Michele Zocchi developed ultrasonographically guided liposuction in the early 1990s, expanding the applicability of the procedure to previously unfavorable, fibrous regions such as the buttocks (1,6). That same decade, David Apfelberg introduced minimally invasive laser-assisted liposuction, which reduced damage to neurovascular structures and promoted tissue tightening for improved aesthetic results (1). More recently, power-assisted liposuction has further expanded the use and popularity of the procedure by reducing operator fatigue during large-volume or fibrous-tissue cases (1,5). Given the historically high complication profile and persistent misconceptions surrounding liposuction, the technique was



Figure 1-Liposuction cannulas



Figure 2- ultrasound assisted liposuction

Liposuction and Body Contouring Surgery: A Comprehensive Review of Techniques, Safety, And Evolving Evidence

not formally taught in United States plastic surgery residency programs until the 1980s; today it ranks as the second most commonly performed cosmetic procedure after breast augmentation among board-certified plastic surgeons (5).

III. ANATOMICAL CONSIDERATIONS

A clear understanding of adipose tissue architecture underlies the safe and effective practice of liposuction. Adipocytes are composed largely of triglycerides that occupy the majority of the cell's volume, displacing the nucleus to one side (5). In the subcutaneous space, fat is organized into three principal layers—superficial (immediately beneath the dermis), middle or intermediate, and deep—with each layer composed of fat lobules separated by fibrous septa that house neurovascular channels (5).

During childhood, weight gain stimulates an increase in the absolute number of adipocytes through differentiation of adipose-derived stem cells. Once adulthood is reached, however, the number of adipocytes becomes essentially fixed, and subsequent weight gain is instead mediated by an increase in the size of existing cells (5). This biological feature explains why liposuction can produce durable contour change: by permanently removing a defined population of fat cells, the procedure targets adipocytes that might otherwise be resistant to diet and exercise (5). Lipoaspiration produces a bimodal reduction in adipocyte number—an immediate reduction from direct mechanical removal, followed by a secondary reduction approximately six to twelve weeks later as damaged but incompletely aspirated cells undergo resorption (5). Nevertheless, because remaining adipocytes retain the capacity to enlarge and adipose-derived stem cells retain the capacity to differentiate, future weight gain can still negate the contour benefits achieved through surgery (1,5).

Fatty and fibrous tissue characteristics vary considerably by anatomical region. In the arms and thighs, fat is loosely organized with minimal fibrous septation between layers, whereas in the back the fatty layers are densely tethered by fibrous attachments that organize the tissue into a compact, structurally rigid system (5). Five recognized zones of adherence—the lateral gluteal depression, gluteal crease,

distal posterior thigh, midmedial thigh, and inferolateral iliotibial tract—represent regions where superficial subcutaneous tissue adheres firmly to the underlying deep fascia (1,5). Because these zones of adherence define the natural

shape and surface topography of the body, aggressive suctioning within them substantially increases the risk of visible contour deformity (1). The most common region for liposuction is the broad area extending from the inframammary fold to the gluteal fold, encompassing the abdomen, flanks, trochanteric region, lumbar region, and buttocks (1).

Traditionally, the middle and deep adipose layers were considered the appropriate targets for liposuction, with the superficial layer left undisturbed because of the perceived risk of contour irregularity (5). Gasparotti subsequently popularized superficial liposuction for selected indications, including treatment of aged or flaccid skin, and contemporary practice now recognizes superficial-layer liposuction—including the deliberate exploitation of vertical fibrous septa to create defined abdominal musculature, known as “abdominal etching”—as a legitimate extension of traditional technique rather than an absolute contraindication (1,5).

For ideal aesthetic outcomes, patients should possess adequate skin elasticity and fall within approximately 20 to 30 percent of their ideal body weight (1). Liposuction is not, and should not be presented as, a weight-loss intervention; rather, it is most appropriately conceived as a body-contouring technique for patients with isolated, disproportionate adipose deposits (5). Patients with a body mass index greater than 30 are generally better served by alternative weight-management strategies, although some authors note that liposuction may, in select circumstances, serve as a motivational adjunct that encourages patients to pursue more comprehensive lifestyle modification (5).

IV. COSMETIC AND NONDOMESTIC INDICATIONS

Cosmetic Applications

Liposuction is used to achieve body contouring through removal of excess, anatomically isolated fat deposits. Common treatment sites include the abdomen, flanks, trochanteric and lumbar regions, buttocks, breasts, thighs, calves, neck, and arms (1,5). Liposuction is increasingly employed as an adjunct to other aesthetic procedures, including breast augmentation, cervicoplasty, abdominoplasty, gluteal fat transfer, and body contouring following bariatric surgery (1). The procedure may also be tailored to achieve gender-specific aesthetic



Figure 3.- Immediate postoperative liposuction




Figure 5.- Hip and glutes contour

Liposuction and Body Contouring Surgery: A Comprehensive Review of Techniques, Safety, And Evolving Evidence

goals: in women, the objective is typically to create shapely contours of the breasts, waist, hips, and buttocks, whereas in men the goal is often to achieve upper-body dominance through reduction of flank adiposity (1).

Gluteal Fat Transfer

Gluteal fat transfer, popularly known as the Brazilian butt lift, applies liposuction principles by harvesting large fat volumes from a donor site, such as the abdomen or inner thighs, for autologous transfer to the buttocks (1). High-volume fat transfer, generally defined as more than 1,000 mL per buttock, has historically carried elevated risk of graft-site infection and donor-site seroma, although newer evidence suggests that high-volume transfer may be performed safely with appropriate technique (1). The procedure's popularity has grown rapidly, driven in part by shifting beauty standards and celebrity visibility: more than 26,000 gluteal fat transfer procedures were performed in 2018 alone, a 16 percent increase over the previous year and a 132 percent increase since 2013 (1). This growth has occurred alongside serious safety concerns. Reports of fatal pulmonary fat embolism following injury to gluteal veins, with an estimated mortality rate of 1 in 3,000, have prompted the formation of the Multi-Society Gluteal Fat Grafting Task Force to investigate and improve patient safety for this specific procedure (1).

Noncosmetic Applications

Liposuction also serves a number of reconstructive and medical indications. These include management of lipomas and angioliomas with minimal scarring; lipedema, in which abnormal subcutaneous fat deposition in the lower limbs impairs mobility; lymphedema refractory to conservative treatment; and lipodystrophy syndromes, whether congenital or acquired, in which autologous fat transfer can restore lost volume in areas such as the feet or buttocks to relieve physical discomfort (1). Additional noncosmetic uses include treatment of cervicodorsal lipodystrophy associated with Cushing syndrome or antiretroviral therapy, correction of gynecomastia in men and macromastia in women in conjunction with mammoplasty, reduction of fat at planned surgical sites (for example, prior to tracheostomy, colostomy, or urostomy) in obese patients, thinning of reconstructive flaps to improve aesthetic outcomes, and harvesting of adipose tissue for “lipofilling” in breast reconstruction, burn management, and scar revision, given the presence of adipose-derived stem cells within harvested fat (1). Liposuction has additionally been described as an effective treatment for axillary hyperhidrosis, in which removal of fat from the axillary vault, combined with curettage against the dermis, reduces the density of eccrine and apocrine glands and offers a durable alternative to repeated botulinum toxin injection (6).

Although no absolute contraindications to liposuction exist, several relative contraindications warrant careful preoperative evaluation. Anticoagulants and medications

that interfere with lidocaine metabolism should be discontinued before surgery, and poor skin elasticity in elderly patients increases the risk of unsatisfactory postoperative skin draping (1). Patients with suspected body dysmorphic disorder may require psychiatric evaluation before proceeding, and those with diabetes mellitus, cardiac disease, or hepatic disease may require medical clearance, with poorly controlled diabetes specifically recognized as a risk factor for postoperative infection (1,5).

V. SURGICAL TECHNIQUES AND ASSISTIVE TECHNOLOGIES

Despite the proliferation of more than fifty liposuction-related devices and technique modifications introduced over the past several decades, traditional suction-assisted lipectomy remains the most widely used approach (1,3,5). All assistive technologies—ultrasound, power, laser, radiofrequency, and water-jet systems—rely on suction as the final common step for fat aspiration, even though their respective acronyms tend to emphasize the energy modality rather than the suction mechanism itself (5).

A. Suction-Assisted Lipectomy

In suction-assisted lipectomy (SAL), negative pressure—typically equivalent to one atmosphere, or approximately 760 mmHg—is applied through a blunt-tipped cannula to remove fat by mechanical avulsion: the suction traps tissue within the cannula tip, and forward-backward cannula motion subsequently tears the tissue free (5). This remains the technique with the lowest equipment cost, the most established safety record, and the most straightforward learning curve, though it is comparatively labor-intensive for the surgeon and offers minimal direct skin contraction (5). Suctioning from the superficial fat layer should generally be avoided to prevent dimpling and contour irregularity, given the vertical fibrous septa contained within this layer, although deliberate, controlled use of this anatomy can be exploited to create defined abdominal contours (1).

B. Ultrasound-Assisted Lipectomy

Ultrasound-assisted lipectomy (UAL), first developed by



Figure 6.- Ultrasound-guided aspiration

Zocchi, uses mechanical and thermal ultrasonic energy to selectively emulsify adipocytes prior to aspiration, a property that makes it particularly advantageous in dense, fibrous regions such as the back, chest, and upper flank (1,5). Device design varies between solid probes,

Liposuction and Body Contouring Surgery: A Comprehensive Review of Techniques, Safety, And Evolving Evidence

which are more efficient at emulsification but require a separate aspiration stage, and hollow probes, which are less efficient but allow single-stage emulsification and aspiration (5). Because of the risk of thermal injury at the skin entry site, protective sheaths and damp sponge barriers are required whenever ultrasound energy is used (5).

The third-generation ultrasound device, marketed as VASER (vibration amplification of sound energy at resonance), incorporates titanium probes with rings or grooves that redistribute energy away from the cannula tip, thereby reducing the complications—seroma, pigmentation change, and thermal injury—that affected earlier-generation devices (5). A consistently reported advantage of ultrasound-assisted lipectomy is reduced intraoperative blood loss attributable to coagulation of adjacent small vessels: one comparative study reported a 7.5-fold reduction in blood loss with UAL relative to SAL (3.98 percent versus 0.61 percent of total lipoaspirate, $p < 0.0001$) (3,5). This blood-sparing effect has been corroborated across multiple independent investigations. In a prospective study of 22 patients undergoing third-generation internal ultrasound-assisted liposuction, hemoglobin declined by a mean of 0.53 ± 0.36 g/dL and hematocrit by 1.50 ± 1.13 percentage points for each liter of aspirate removed, both significant reductions from preoperative baseline, but smaller in magnitude than historically reported with SAL (4). In that same study, the amount of aspirated supernatant accounted for 44.4 percent of the variance in hemoglobin change and 30.9 percent of the variance in hematocrit change, and a strong positive correlation was observed between greater hemoglobin and hematocrit reduction and both female sex and smoking status (4).

A separate comparative trial evaluating VASER, a proprietary device termed LipoSaver2000, and traditional SAL found no significant differences among the three modalities in hemoglobin or hematocrit content of the lipoaspirate, although aspirate volume was significantly higher in the SAL group than in the VASER group ($p = 0.001$) (2). This finding stands in contrast to several earlier reports describing a clear ultrasound-related blood-sparing benefit, illustrating that the magnitude of this effect may vary by device, technique, and patient population (2).

Notwithstanding these blood-loss advantages, a comprehensive evidence-based review identified only five studies meeting strict methodological criteria for direct comparison of UAL and SAL, and concluded that despite well-documented clinical efficiency, relative safety, and reduced rates of postoperative anemia, there remains insufficient long-term data to confirm that UAL produces superior aesthetic outcomes or skin tightening compared with SAL (3). The same review noted that UAL carries a higher overall complication rate (35.2 percent) than other fat-removal methods and recommended that the technology be reserved for patients with low body mass index requiring low-to-moderate volume fat removal, cautioning that



Figure 7.- Radiofrequency-Assisted Liposuction.

offering UAL to patients with excessive fat or skin laxity for whom more invasive surgery is indicated may be misleading regarding achievable outcomes (3).

C. Power-Assisted Lipectomy

Power-assisted

lipectomy (PAL) incorporates an electric or pneumatic mechanism that drives rapid reciprocating motion of the cannula tip, reducing the physical effort required of the surgeon and improving economy of motion, particularly in fibrous regions such as the male breast or in revision cases (5). Reported increases in fat extraction efficiency with powered devices, relative to manual technique, have reached as high as 45 percent (5). The principal disadvantage of PAL is the vibration transmitted to the surgeon's hand, along with operative noise, both of which become more burdensome during lengthy large-volume procedures (5).

D. Laser-Assisted Lipectomy

Laser-assisted lipectomy (LAL) employs a laser fiber introduced through a small incision to rupture adipocyte membranes, releasing cellular contents into the extracellular space prior to aspiration (1,5). The accompanying thermal coagulation of small vessels and collagen fibers is associated with decreased intraoperative blood loss and improved postoperative skin tightening, though at the cost of more expensive equipment and prolonged procedure time (5). Complication rates with LAL are reported to be low, with one study finding a complication rate of 0.93 percent, including skin burns and localized infection (1).

E. Radiofrequency- and Water-Assisted Lipectomy

Radiofrequency-assisted lipectomy delivers energy simultaneously to the dermis and underlying subcutaneous tissue through a handpiece containing both external and internal probes, with continuous temperature monitoring designed to prevent thermal overshoot; one study found average linear skin contraction of 31 percent with this technology, exceeding that reported for other liposuction modalities, theoretically attributable to collagen fiber contraction and neocollagen formation (5). Water-assisted lipectomy uses a fanning jet of pressurized tumescent fluid to dissect and loosen fat for simultaneous aspiration through a secondary channel within the cannula; this remains the most recently introduced and least-studied of the assistive technologies, and further comparative data are needed before firm conclusions can be drawn regarding its relative efficacy (5).

VI. ANESTHETIC PRINCIPLES AND FLUID MANAGEMENT

The introduction of tumescent anesthesia by Klein in 1987 stands as the single most important advance in the modern practice of liposuction, eliminating the requirement for general anesthesia in many cases, reducing blood loss and associated transfusion needs, and shortening overall patient recovery time (5,6). Tumescent solution typically combines a dilute lidocaine concentration (0.05 to 0.1 percent), epinephrine at a dilution of approximately 1:1,000,000, and sodium bicarbonate buffer within a balanced crystalloid carrier such as lactated Ringer's solution or normal saline (6). The so-called "superwet" or tumescent infiltration techniques use a defined ratio of infiltrate to anticipated aspirate and are associated with a blood content of less than 2 percent of total fluid removed, a marked improvement over the 30 to 45 percent blood content historically observed with dry-technique liposuction (5).

Tumescent anesthesia permits substantially higher lidocaine dosing than conventional subcutaneous infiltration. Whereas concentrated subcutaneous lidocaine is limited to a maximum of 5 to 7 mg/kg, large-volume tumescent infiltration has been demonstrated to be safe at doses up to 55 mg/kg, although the American Society of Plastic Surgeons continues to recommend a more conservative maximum of 35 mg/kg, based on Klein's original 1990 publication (5). This favorable safety margin results from several pharmacokinetic factors: lidocaine is lipophilic and becomes sequestered within fat cells, producing a slow release over the following 12 to 24 hours, with lidocaine and its active metabolite reaching peak plasma concentrations between 8 and 18 hours after injection; in addition, much of the infiltrated solution is subsequently removed through aspiration before full systemic absorption can occur (5,6). Bupivacaine is specifically discouraged as a component of tumescent solution because of its longer duration of action, slower elimination, and greater potential for cardiac, neurologic, and hematologic toxicity (5).

Detumescence the process by which infiltrated fluid spreads through tissue planes and exerts its full pharmacologic effect generally requires 20 to 30 minutes, although many surgeons proceed after approximately 10 to 15 minutes in order to preserve operative efficiency, particularly when tissue blanching or turgor provides an adequate clinical endpoint (5,6). For patients undergoing large-volume liposuction, defined as more than 4 liters of aspirate, additional intravenous crystalloid resuscitation approximately 1 mL for every 4 mL of lipoaspirate exceeding 4,000 mL is recommended, in addition to replacement of any preoperative fluid deficit and routine maintenance fluids (5). Hemodynamic studies have demonstrated that large-volume liposuction is associated with measurable physiologic changes, including increased heart rate, mean pulmonary artery pressure, cardiac index,

and stroke volume index, alongside decreased mean arterial pressure and systemic vascular resistance index, with plasma epinephrine concentrations peaking five to six hours after induction of anesthesia (5). More recent data indicate that the risk of liposuction-related complications correlates more closely with body mass index than with absolute aspirate volume, with a proposed safe upper limit of approximately 100 mL of aspirate per unit of body mass index (5).

VII. BLOOD LOSS AND THE ROLE OF TRANEXAMIC ACID

Blood loss has long been recognized as one of the principal limiting factors in large-volume liposuction, constraining the amount of fat that can safely be aspirated in a single session (2,4). Approximately 20 to 45 percent of lipoaspirate volume is blood when liposuction is performed without fluid infiltration; this proportion falls to roughly 10 to 30 percent with the wet technique, to approximately 15 percent with the addition of epinephrine, and to as low as 2 percent of aspirated volume with the superwet technique (4,5). Aspiration volume, operative duration, and patient body weight have been identified as the most important determinants of overall blood loss (4).

Tranexamic acid (TXA), an antifibrinolytic agent with established efficacy in reducing blood loss in cardiac, obstetric, trauma, and orthopedic surgery, has more recently been investigated as an adjunct in plastic and aesthetic surgery, including rhinoplasty, facelift, breast reduction, and dermatologic procedures (2). A 2016 meta-analysis reported that TXA could reduce blood loss and transfusion requirements in craniomaxillofacial and plastic surgery generally (2), and individual studies in rhinoplasty and dermatologic surgery have similarly reported reduced intraoperative bleeding and improved surgical field visibility with TXA use (2).

Evidence specific to liposuction, however, remains comparatively limited and inconsistent. Cansancao and colleagues, in a study of ten women, reported that preoperative and postoperative intravenous administration of 10 mg/kg TXA significantly reduced blood loss per liter of lipoaspirate, finding 8.8 mL of blood loss per liter of supernatant in the TXA-treated group compared with 20.1 mL in the untreated group—a 56.2 percent relative reduction—along with a 48 percent reduction in hematocrit decline (2,4). Rodriguez-Garcia and colleagues, studying 50 women who received 1,000 mg of TXA added to tumescent solution, similarly reported significantly lower hematocrit decline relative to aspirated volume in the TXA group compared with controls (2). In a study of 36 patients undergoing combined liposuction and resection for breast reduction, infiltration of 0.5 g/5 mL TXA solution into one breast, with a TXA-free solution infiltrated into the contralateral breast, resulted in 38 percent less blood volume in the lipoaspirate from the TXA-treated side (2). El Minawi

and colleagues compared locally infiltrated TXA (8.1 mg/kg) with intravenously administered TXA (7.8 mg/kg) and found that local infiltration significantly reduced blood loss per milliliter of aspirate and lowered hematocrit values in the lipoaspirate, whereas intravenous administration produced only a numerical, non-significant decrease (2).

In direct contrast to these findings, a prospective study comparing VASER, LipoSaver2000, and traditional SAL techniques in which 500 mg of TXA was added to 1,000 cc of tumescent solution and applied to one flank, with the contralateral flank receiving tumescent solution without TXA found no significant difference in hemoglobin or hematocrit content of the lipoaspirate between TXA-treated and untreated sides, regardless of the liposuction technique employed (2). The authors of that study attributed this discrepancy to the comparatively lower TXA dose used (500 mg per 1,000 cc of tumescent solution) relative to the higher doses reported in studies demonstrating a positive effect, suggesting that the concentration of TXA within the infiltration solution may be an important determinant of its hemostatic efficacy in this setting (2). The authors further cautioned that, irrespective of TXA's apparent lack of effect on bleeding in their cohort, incorrect or overly superficial infiltration technique carries an independent risk of superficial skin necrosis through injury to the dermal and subdermal vascular network (2).

Taken together, the available evidence suggests that TXA may offer a meaningful blood-sparing benefit in liposuction when administered at sufficiently high concentrations, whether by local infiltration or intravenous route, but that this benefit is not uniformly observed across all reported protocols. The heterogeneity in dosing, administration route, and technique across published series limits the ability to draw firm, generalizable conclusions, and several authors have called explicitly for larger, standardized trials before TXA can be considered a routine component of liposuction practice (2).

VIII. COMPLICATIONS AND SAFETY OUTCOMES

Liposuction is associated with a comparatively low overall complication rate relative to other surgical procedures. One large series reported an overall complication rate of 2.4 percent, rising to 3.5 percent when liposuction was combined with other procedures, but falling to only 0.7 percent when liposuction was performed as a solitary procedure (1). A separate analysis identified an overall contour-irregularity incidence of 2.7 percent, with hematoma occurring in 0.15 percent of cases and deep venous thrombosis or pulmonary embolism in 0.06 percent (5).

Contour deformity is consistently identified as the most common complication of liposuction, affecting as many as 9 percent of patients in the form of soft-tissue depressions, elevations, panniculus folds, or wrinkling (1). Strategies to reduce this risk include the use of smaller-diameter

cannulas, avoidance of superficial-layer suctioning outside of deliberate etching techniques, employment of a criss-crossing cannula pattern, and deliberate slight undercorrection to account for postoperative fat lysis (1). Seroma and hematoma, while uncommon, may arise from blind cannula injury to small perforating vessels or lymphatic channels; the use of progressive tension sutures, a technique borrowed from abdominoplasty closure, has been shown to reduce seroma rates from 9 percent to 2 percent (1).

Wound infection occurs in fewer than 3 percent of inpatient liposuction cases and in approximately 1 percent of outpatient procedures, with low rates generally attributable to surgeon expertise, prophylactic antibiotic use, and sterile technique (1). Although uncommon, early superficial infection can progress to more severe presentations such as necrotizing fasciitis or toxic shock syndrome, both of which constitute surgical emergencies (1,5).

Fat embolism, although rare, carries a mortality rate of 10 to 15 percent and most often manifests within 12 to 72 hours after surgery, warranting vigilant postoperative monitoring (1). One 2018 study estimated that at least 17 patients worldwide had experienced clinically significant fat embolization following liposuction with or without subsequent fat grafting, while more recent estimates suggest that the true global figure—particularly when gluteal fat grafting is included—may exceed 135 cases (1). Gluteal fat transfer in particular is recognized as carrying the highest mortality rate of any aesthetic surgical procedure, a risk specifically related to fat embolism following inadvertent injury to gluteal veins (1). By contrast, the overall incidence of venous thromboembolic events after conventional liposuction—encompassing both deep venous thrombosis and pulmonary embolism—is low, at approximately 0.03 percent, although pulmonary embolism remains the most common cause of death after the procedure, contributing to an overall mortality rate of approximately 0.01 percent (1). Same-day ambulation following surgery is encouraged specifically to reduce thromboembolic risk (1).

Systemic complications that may arise in the weeks to months following surgery include edema, lymphedema, wound dehiscence, hypertrophic scarring, ecchymosis, and skin laxity (1). Blind cannula injury carries the potential, albeit rare, for abdominal wall injury, bowel perforation, or vascular injury, and inexperienced use of ultrasound-assisted devices in particular has been associated with abdominal perforation, in part because the thermal and mechanical properties of ultrasound energy reduce the surgeon's tactile feedback during cannula passage (1,6). Skin devascularization and necrosis may occur if suctioning is performed too close to the dermal undersurface, injuring the dermal vascular plexus (1). Breast augmentation performed with autologous fat transfer may produce fat necrosis that mimics microcalcifications suspicious for malignancy on subsequent mammographic imaging; reassuringly, the

Liposuction and Body Contouring Surgery: A Comprehensive Review of Techniques, Safety, And Evolving Evidence

incidence of such imaging findings appears similar to that observed in patients who have not undergone fat transfer, indicating that the procedure does not meaningfully impair breast cancer detection (1).

Risk factors that increase the likelihood of surgical complications include cardiovascular disease, pulmonary disease, diabetes, vascular disease, tobacco use, active or recent infection near the planned surgical site, and a prior history of venous thromboembolism (1). As with any surgical procedure, liposuction produces a transient elevation in acute inflammatory markers such as interleukin-6 and C-reactive protein, though without evidence of progression to renal disease or chronic systemic inflammation; indeed, some studies suggest that permanent adipocyte removal may confer long-term metabolic benefits, including improved insulin sensitivity and reduced systemic inflammation, although this remains an area of active investigation with conflicting results across studies (1).

Large-volume and debulking liposuction, frequently employed in patients with lipedema or massive weight loss, presents a distinct complication profile. Current recommendations call for hospitalization when aspirate volumes exceed 5,000 mL, given the potential for significant postoperative blood loss even when the aspirate itself does not appear grossly bloody (5). Postoperative edema in such cases can be substantial and difficult to manage with standard compression garments, occasionally progressing to cellulitis or necessitating lymphedema-grade compression therapy (5).

IX. FUTURE DIRECTIONS

Liposuction continues to evolve along several parallel trajectories. Noncosmetic indications are expanding, particularly the use of fat grafting for breast, facial, and pedal reconstruction, even though liposuction performed for functional or reconstructive purposes is rarely covered by Medicare or third-party insurance, even in cases involving genuine functional impairment (1). Investigation into the long-term metabolic consequences of large-volume fat removal, including effects on insulin sensitivity and cardiovascular risk factors, remains ongoing and has produced conflicting results to date (1).

Considerable research interest has also focused on noninvasive alternatives to surgical liposuction, including cryolipolysis (marketed as CoolSculpting), deoxycholic acid injection (marketed as Kybella), and radiofrequency-based skin tightening (marketed as Thermage); while these modalities address some of the same patient concerns around localized adiposity and skin laxity without surgical scarring, they remain distinct from, and are not a substitute for, surgical liposuction in patients with substantial fat or skin excess (1).

Within the surgical domain, evidence-based scrutiny of energy-based devices is likely to intensify. Several reviewers have cautioned that despite considerable industry

investment and marketing around technologies such as VASER-assisted high-definition liposculpture, rigorous long-term comparative data confirming superior aesthetic outcomes or skin retraction relative to traditional SAL remain limited (3). Combined approaches—for example, pairing VASER with helium-based plasma radiofrequency technology, or supplementing high-definition liposculpture with ultrasound-guided rectus fat grafting techniques for abdominal etching—represent active areas of innovation, though objective, standardized outcome measures are still needed to validate their comparative benefit (3). Similarly, continued refinement of pharmacologic adjuncts such as tranexamic acid, particularly with respect to optimal dosing and route of administration, may further improve the safety profile of large-volume procedures (2). As the field progresses, the central challenge will be balancing genuine technological advancement against marketing-driven adoption, ensuring that new devices and adjuncts are incorporated into practice only as supporting evidence warrants (3,5).

CONCLUSIONS

Liposuction has evolved from a hazardous, poorly understood procedure in the early twentieth century into one of the safest and most frequently performed surgical interventions in contemporary aesthetic and reconstructive practice. The introduction of tumescent anesthesia fundamentally transformed the safety profile of the procedure, and subsequent technological refinements—ultrasound, power, laser, and radiofrequency assistance—have each sought to address specific limitations of traditional suction-assisted technique, with varying degrees of supporting evidence. Comparative data consistently support a blood-sparing advantage for ultrasound-assisted liposuction, although claims of superior long-term skin tightening remain insufficiently substantiated by rigorous, controlled, long-term studies. Pharmacologic strategies such as tranexamic acid show promise in reducing intraoperative blood loss but require further standardization before routine adoption can be recommended. Across all techniques, contour deformity remains the most common complication, while gluteal fat transfer carries a disproportionately elevated mortality risk that warrants continued safety investigation through dedicated multi-society task forces. As both cosmetic demand and reconstructive applications for liposuction continue to expand, ongoing evidence-based evaluation—rather than reliance on marketing claims for newer technologies—should guide the selection of technique for individual patients

REFERENCES

1. Wu S, Coombs DM, Gurunian R. Liposuction: Concepts, safety, and techniques in body-contouring surgery. *Cleve Clin J Med*. 2020;87(6):367-375. doi:10.3949/ccjm.87a.19097.

Liposuction and Body Contouring Surgery: A Comprehensive Review of Techniques, Safety, And Evolving Evidence

- II. Kandulu H. Comparative analysis of the effectiveness of tranexamic acid in controlling surgical blood loss in patients undergoing liposuction. *Aesthetic Plast Surg.* 2025;49:4343-4349. doi:10.1007/s00266-025-04852-6.
- III. Atieh B, Baajour J, El Merkabaoui H, Makkawi K. Literature review and evidence-based comparative analysis of traditional suction-assisted lipectomy (SAL) vs. ultrasound-assisted liposuction (UAL) reported outcomes. *Aesthetic Plast Surg.* 2025;49:4648-4655. doi:10.1007/s00266-025-04872-2.
- IV. Kandulu H. Evaluation of blood loss in third-generation internal ultrasound-assisted liposuction. *Aesthetic Plast Surg.* 2024;48:152-156. doi:10.1007/s00266-023-03633-3.
- V. Beidas OE, Gusenoff JA. Update on liposuction: what all plastic surgeons should know. *Plast Reconstr Surg.* 2021;147(4):658e-668e. doi:10.1097/PRS.00000000000007419.
- VI. Coleman WP, Hendry SL. Principles of liposuction. *Semin Cutan Med Surg.* 2006;25:138-144. doi:10.1016/j.sder.2006.06.006.