

Management of the Difficult Airway In Anesthesia: Prediction, Oxygenation, and Rescue Strategies

Adela Angélica Córdova Fernandez

Hospital de Gineco-Obstetricia IMSS No. 4 “Dr. Luis Castelazo Ayala”, Mexico City, Mexico

ABSTRACT

Difficult airway management remains one of the most critical areas in anesthetic practice because failure to oxygenate or intubate can rapidly lead to hypoxic brain injury, cardiac arrest, or death. A difficult airway includes situations in which a trained clinician encounters difficulty with facemask ventilation, supraglottic airway placement, laryngoscopy, tracheal intubation, or emergency surgical access. Preoperative assessment can identify high-risk patients but has limited ability to reliably exclude difficulty, making universal preparation essential. Risk factors include high Mallampati score, reduced mouth opening, short thyromental distance, limited neck extension, obesity, obstructive sleep apnea, craniofacial abnormalities, airway masses, trismus, and subglottic stenosis. Contemporary management emphasizes preoxygenation, continuous oxygen delivery, early use of videolaryngoscopy or flexible bronchoscopy when indicated, limitation of repeated attempts, rapid transition between airway devices, mandatory waveform capnography, and timely front-of-neck access when ventilation fails. Awake tracheal intubation remains the preferred strategy for anticipated difficult airway when aspiration risk, rapid desaturation, or impossible rescue ventilation is expected. Successful management depends less on a single device and more on structured decision-making, team communication, oxygenation preservation, and early rescue escalation.

KEYWORDS: difficult airway; anesthesia; tracheal intubation; videolaryngoscopy; awake intubation; supraglottic airway; front-of-neck access; oxygenation..

ARTICLE DETAILS

Published On:
14 October 2026

Available on:
<https://ijmscrs.com>

INTRODUCTION

The difficult airway is a low-frequency but high-consequence event in anesthesia. It may involve difficult facemask ventilation, difficult supraglottic airway placement, difficult laryngoscopy, difficult tracheal intubation, or complete failure of oxygenation. Although impossible intubation and ventilation are rare, they remain among the most feared anesthetic emergencies because neurological injury and death can occur within minutes if oxygenation is not restored.

Modern difficult airway management has shifted from prediction alone toward preparation and rescue planning. Bedside airway tests are useful, but none is sufficiently sensitive or specific to guarantee a safe airway. The uploaded synthesis defines difficult airway as difficulty with facemask ventilation, supraglottic airway placement, laryngoscopy, or tracheal intubation, and notes that impossible intubation-and-ventilation scenarios remain a major contributor to anesthesia-related hypoxic morbidity and mortality.



Intraoperative positioning. Patient in the supine position under general anesthesia, with cervical hyperextension for adequate exposure of the submental and anterior cervical regions.



Example of Hard Airway

Preoperative Prediction and Risk Stratification

Preoperative airway evaluation remains essential, but it should be interpreted as risk stratification rather than certainty. Mallampati class, interincisor distance, thyromental distance, mandibular protrusion, neck mobility, cervical circumference, body habitus, dentition, and history of prior airway difficulty should all be considered together. Multivariate assessment is more useful than any isolated bedside test, particularly because difficult laryngoscopy, difficult ventilation, and difficult rescue oxygenation do not always share the same predictors.

The most important adult predictors include Mallampati III–IV, limited mouth opening, reduced thyromental distance, restricted neck extension, obesity, and obstructive sleep

apnea. Obstructive sleep apnea is especially relevant because it increases the risk of both difficult intubation and difficult ventilation. In children, syndromic craniofacial hypoplasia, micrognathia, and congenital airway abnormalities substantially increase the risk of critical respiratory events. The source also notes that emerging deep learning models have shown diagnostic performance with areas under the receiver operating curve around 0.80–0.84, but these tools should complement rather than replace clinical judgment.

Principles of Management

The central priority in difficult airway management is oxygenation, not intubation. A well-executed airway plan begins before induction with patient positioning, preoxygenation, suction, availability of videolaryngoscope, bougie, supraglottic airway, flexible bronchoscope, front-of-neck access equipment, and a clear plan for failure. The team

Management of the Difficult Airway In Anesthesia: Prediction, Oxygenation, and Rescue Strategies

should know when to stop, when to change devices, when to wake the patient, and when to proceed to emergency surgical access.

Current guidelines emphasize continuous oxygen delivery during airway manipulation and extubation. High-flow nasal oxygen may prolong safe apnea time in high-risk patients, although it does not eliminate the need for decisive action. Waveform capnography is mandatory after intubation because unrecognized esophageal intubation remains a recurring cause of perioperative cardiac arrest. Repeated attempts are dangerous because they produce edema, bleeding, trauma, laryngospasm, and progressive loss of rescue options. The source summarizes guideline recommendations to restrict attempts, call for help early, and transition promptly among tracheal tube, supraglottic airway, and facemask ventilation.

Anticipated Difficult Airway

When difficulty is expected, the decision is whether the airway should be secured awake or after induction. Awake tracheal intubation is favored when loss of spontaneous ventilation would be dangerous, especially in patients with airway obstruction, high aspiration risk, unstable cervical spine, anticipated difficult ventilation, or rapid desaturation risk. Flexible bronchoscopic intubation remains a classic approach, but awake videolaryngoscopy is increasingly used when anatomy permits.

Awake intubation is not automatically safe. Excess sedation can cause hypoventilation, airway obstruction, or hypoxemia. The uploaded synthesis notes that observational registries have reported hypoxemia in up to 50% of awake intubations, often related to sedation-induced hypoventilation. hard airway in anesthesia Therefore, topical anesthesia, minimal titrated sedation, oxygen supplementation, patient cooperation, and immediate rescue readiness are essential.

Unanticipated Difficult Airway

Unanticipated difficulty after induction requires rapid algorithmic progression. Initial optimization includes repositioning, external laryngeal manipulation, suction, change of blade or device, use of bougie, and calling for help. Videolaryngoscopy has become central because it improves glottic visualization, but better visualization does not always guarantee tube delivery. Hyperangulated blades often require a rigid stylet, deliberate tube shaping, and controlled advancement.

If intubation fails, the clinician must return to oxygenation. Facemask ventilation, two-person technique, oral or nasal airway placement, and supraglottic airway insertion should be attempted early rather than after multiple traumatic laryngoscopies. A supraglottic airway may serve as a rescue oxygenation device, a conduit for fiberoptic intubation, or a bridge to awakening the patient. If oxygenation is restored, the safest option may be to wake the patient and replan rather than continue escalating attempts.

Cannot Intubate, Cannot Oxygenate

The “cannot intubate, cannot oxygenate” situation is the critical endpoint of failed airway management. Once facemask ventilation, supraglottic airway ventilation, and tracheal intubation have failed, front-of-neck access should not be delayed. Emergency cricothyrotomy or surgical airway is indicated before severe hypoxemia progresses to cardiac arrest. The source specifically identifies complete failure of mask, supraglottic, and tube ventilation as the trigger for immediate emergency front-of-neck access.

The key error in this scenario is fixation on intubation. Persistent laryngoscopy attempts can convert a partially manageable airway into a bloody, swollen, impossible airway. Team language should be explicit: failed oxygenation should trigger a shared declaration and immediate movement to front-of-neck access.

Pediatric, Obstetric, and High-Risk Groups

Pediatric airway management differs because infants and children desaturate rapidly, have smaller airway diameter, higher oxygen consumption, and age-specific anatomy. Craniofacial syndromes, micrognathia, and low functional reserve increase risk. Pediatric guidelines favor age-adapted equipment, appropriate videolaryngoscope blades, adequate depth of anesthesia, and careful use of neuromuscular blockade when indicated.

Obstetric airway management is uniquely high risk because pregnancy combines airway edema, aspiration risk, reduced functional residual capacity, and rapid desaturation. Anticipated difficult obstetric airway often favors awake intubation, while failed intubation after induction requires early consideration of supraglottic rescue, awakening, or proceeding only if maternal or fetal urgency demands it.

Patients with obesity and obstructive sleep apnea require particular attention to positioning, ramped alignment, preoxygenation, apneic oxygenation, and postoperative airway safety. In these patients, the airway plan does not end after intubation; extubation and recovery may be equally dangerous.

Extubation Strategy

Extubation is part of difficult airway management. A patient who was difficult to intubate may be difficult to reintubate, especially after airway edema, bleeding, prolonged surgery, cervical fixation, or maxillofacial procedures. Extubation should occur only when oxygenation, ventilation, neuromuscular recovery, hemodynamics, and airway patency are adequate. In high-risk patients, extubation over an airway exchange catheter, delayed extubation in a monitored setting, or staged extubation may be appropriate.

DISCUSSION

The main lesson from modern difficult airway literature is that prediction is imperfect. A reassuring airway examination does not eliminate risk, and an abnormal airway examination does not by itself dictate one technique. The safest approach combines risk recognition with universal readiness. Every anesthetic should have a primary plan, backup oxygenation

Management of the Difficult Airway In Anesthesia: Prediction, Oxygenation, and Rescue Strategies

strategy, rescue supraglottic option, and front-of-neck access pathway.

Videolaryngoscopy, flexible bronchoscopy, high-flow oxygen, and supraglottic devices have improved airway management, but devices do not replace judgment. The clinician must avoid repeated attempts, recognize failure early, and prioritize oxygenation over procedural persistence. Awake intubation remains essential when rescue after

induction is unlikely, but it must be performed with careful topicalization, minimal sedation, and continuous monitoring. The most dangerous airway is not only the anatomically difficult airway, but the airway managed without a clear stopping rule. Structured algorithms, team communication, capnography, and early escalation remain the foundation of safe practice.



Figure 1. Mallampati score

CONCLUSION

Difficult airway management in anesthesia requires preparation, oxygenation preservation, limited attempts, and rapid rescue escalation. Preoperative assessment identifies high-risk patients but cannot reliably exclude difficult intubation or ventilation. Awake tracheal intubation is preferred when induction would create unacceptable risk, while unanticipated difficulty should be managed through an

algorithm that prioritizes oxygenation over repeated laryngoscopy. Videolaryngoscopy, supraglottic airway devices, flexible bronchoscopy, high-flow oxygen, and capnography are essential tools, but successful outcomes depend on timely decision-making. Complete failure of facemask, supraglottic, and tracheal oxygenation requires immediate front-of-neck access. The goal is not simply to place a tube, but to prevent hypoxia.

Clinical situation	Preferred strategy	Key risk to avoid
Predicted difficult intubation with preserved ventilation	Videolaryngoscopy or flexible bronchoscopic plan with backup devices ready	Assuming prediction is perfect
Predicted difficult intubation with aspiration or rapid desaturation risk	Awake tracheal intubation with topical anesthesia and spontaneous ventilation	Oversedation and hypoventilation
Failed first intubation attempt	Reoxygenate, optimize position, change device or operator	Repeating the same failed maneuver
Failed intubation but ventilation possible	Use supraglottic airway or mask ventilation; consider waking the patient	Fixation on intubation
Failed intubation and failed oxygenation	Immediate emergency front-of-neck access	Delayed cricothyrotomy
High-risk extubation	Planned staged extubation or monitored delayed extubation	Treating extubation as routine

REFERENCES

- I. Apfelbaum JL, Hagberg CA, Connis RT, Abdelmalak B, Agarkar M, Dutton RP, et al. 2022 American Society of Anesthesiologists Practice Guidelines for Management of the Difficult Airway. *Anesthesiology*. 2022. doi:10.1097/ALN.0000000000004002
- II. Rosenblatt WH, Yanez N. A decision tree approach to airway management pathways in the 2022 Difficult Airway Algorithm of the American Society of Anesthesiologists. *Anesth Analg*. 2022;134:910-915. doi:10.1213/ANE.0000000000005930
- III. Xia M, Zuo M, Deng XM, Xue FS, Battaglini D, Aggarwal V, et al. Expert consensus on difficult

Management of the Difficult Airway In Anesthesia: Prediction, Oxygenation, and Rescue Strategies

- airway assessment. *Hepatobiliary Surg Nutr.* 2023;12:545-566. doi:10.21037/hbsn-23-46
- IV. Chen H, Zheng Y, Fu Q, Li P. A review of the current status and progress in difficult airway assessment research. *Eur J Med Res.* 2024;29. doi:10.1186/s40001-024-01759-x
- V. Seet E, Nagappa M, Wong DT. Airway management in surgical patients with obstructive sleep apnea. *Anesth Analg.* 2021;132:1321-1327. doi:10.1213/ANE.0000000000005298
- VI. Heidegger T, Pandit JJ. Difficult airway management: from the power of prediction to the art of management. *Anesth Analg.* 2025;140:290-294. doi:10.1213/ANE.0000000000007202
- VII. Myatra SN, Shah AP, Ramkumar V, Kundra P, Patwa A, Shetty S, et al. All India Difficult Airway Association 2025 Guidelines for the management of unanticipated difficult airway in adults under general anaesthesia. *Indian J Anaesth.* 2025;69:1117-1141. doi:10.4103/ija.ija_1210_25
- VIII. Biebuyck JF, Benumof JL. Management of the difficult adult airway with special emphasis on awake tracheal intubation. *Anesthesiology.* 1991;75:1087-1110. doi:10.1097/00000542-199112000-00021
- IX. Yang IT, Tung A, Flores KS, Berenhaut K, Choi JA, Bryan Y. Clinical decision-making and process complications during anticipated difficult airway management for elective surgery. *Anesth Analg.* 2024;140:295-305. doi:10.1213/ANE.0000000000007049
- X. Disma N, Asai T, Cools E, Cronin A, Engelhardt T, Fiadjoe J, et al. Airway management in neonates and infants: European Society of Anaesthesiology and Intensive Care and British Journal of Anaesthesia joint guidelines. *Br J Anaesth.* 2023. doi:10.1016/j.bja.2023.08.040
- XI. Engelhardt T, Virág K, Veyckemans F, Habre W. Airway management in paediatric anaesthesia in Europe—insights from APRICOT. *Br J Anaesth.* 2018;121:66-75. doi:10.1016/j.bja.2018.04.013
- XII. Kojima T, Yamauchi Y, Watanabe F, Ichiyanagi S, Kobayashi Y, Yu KH, et al. Adverse events associated with airway management in pediatric anesthesia: J-PEDIA study. *Anesthesiology.* 2025;143:835-850. doi:10.1097/ALN.0000000000005646
- XIII. Mushambi MC, Athanassoglou V, Kinsella S. Anticipated difficult airway during obstetric general anaesthesia: narrative literature review and management recommendations. *Anaesthesia.* 2020;75. doi:10.1111/anae.15007
- XIV. Cook TM, Oglesby F, Kane A, Armstrong R, Kursumovic E, Soar J. Airway and respiratory complications during anaesthesia and associated with perioperative cardiac arrest as reported to the 7th National Audit Project. *Anaesthesia.* 2023;79. doi:10.1111/anae.16187
- XV. Peterson GN, Domino KB, Caplan RA, Posner KL, Lee LA, Cheney FW. Management of the difficult airway: a closed claims analysis. *Anesthesiology.* 2005;103:33-39. doi:10.1097/00000542-200507000-00009